

cITP

panzyga®

Immune Globulin
Intravenous (Human) - ifas
10% Liquid Preparation

Infusion Rate Chart for Chronic Immune Thrombocytopenia (cITP)

Use this Panzyga Infusion Rate Chart to calculate the appropriate infusion rate for your adult patients with cITP.

- 2 g/kg, divided into 2 daily doses of 1 g/kg (10 mL/kg) given on 2 consecutive days.¹
- The initial infusion rate should be maintained for 30 minutes. Following the initial infusion, and if tolerated, the infusion rate may be gradually increased every 15-30 minutes, as tolerated, to the maximum infusion rate shown in the table below.¹

INDICATION AND USAGE

Panzyga (Immune Globulin Intravenous [Human] - ifas) is indicated for the treatment of primary humoral immunodeficiency (PI) in patients 2 years of age and older; this includes, but is not limited to, congenital agammaglobulinemia, common variable immunodeficiency, X-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies; chronic immune thrombocytopenia (cITP) in adults to raise platelet counts to control or prevent bleeding; and chronic inflammatory demyelinating polyneuropathy (CIDP) in adults to improve neuromuscular disability and impairment.

Patient Weight		mL/hr		
In kg	In lb	First 30 minutes (0.01 mL/kg/min)	Next 15-30 minutes if previous rate tolerated (0.04 mL/kg/min)	Maximum if previous rate tolerated (0.08 mL/kg/min)
40	88	24	96	192
45	99	27	108	216
50	110	30	120	240
55	121	33	132	264
60	132	36	144	288
65	143	39	156	312
70	154	42	168	336
75	165	45	180	360
80	176	48	192	384
85	187	51	204	408
90	198	54	216	432
95	209	57	228	456
100	220	60	240	480
105	231	63	252	504
110	242	66	264	528
115	253	69	276	552
120	264	72	288	576
125	276	75	300	600
130	287	78	312	624
135	298	81	324	648
140	309	84	336	672

Please click [here](#) for Full Prescribing Information, including BOXED WARNING.

octapharma

panzyga®

Immune Globulin
Intravenous (Human) - ifas
10% Liquid Preparation

Contact Octapharma

Octapharma USA, Inc.
117 W. Century Road
Paramus, NJ 07652
Tel: 201-604-1130

Reimbursement Assistance
usreimbursement@octapharma.com
Tel: 800-554-4440

Medical Affairs
usmedicalaffairs@octapharma.com

For all inquiries relating to drug safety, or to report adverse events, please contact our local Drug Safety Officer:
Tel: 201-604-1137
Cell: 201-772-4546
Fax: 201-604-1141
or contact the FDA at 1-800-FDA-1088
or www.fda.gov/medwatch

Reference: 1. Panzyga. Prescribing information. Octapharma USA Inc.; 2021.

© 2025 Octapharma USA, Inc. All rights reserved.
Date of preparation: 05/2025. PANZ-0855

octapharma

IMPORTANT SAFETY INFORMATION (continued)

WARNING: THROMBOSIS, RENAL DYSFUNCTION, AND ACUTE RENAL FAILURE

- **Thrombosis** may occur with immune globulin intravenous (IGIV) products, including Panzyga. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling central vascular catheters, hyperviscosity, and cardiovascular risk factors. Thrombosis may occur in the absence of known risk factors.
- **Renal dysfunction, acute renal failure, osmotic nephrosis, and death** may occur in predisposed patients who receive IGIV products, including Panzyga. Patients predisposed to renal dysfunction include those with a degree of pre-existing renal insufficiency, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs. Renal dysfunction and acute renal failure occur more commonly in patients receiving IGIV products containing sucrose. Panzyga does not contain sucrose.
- **For patients at risk of thrombosis, renal dysfunction, or acute renal failure, administer Panzyga at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity. [see Full Prescribing Information, Warnings and Precautions (5.2, 5.4)]**

Contraindications

PANZYGA is contraindicated in patients who have a history of severe systemic hypersensitivity reactions, such as anaphylaxis, to human immunoglobulin and in IgA-deficient patients with antibodies against IgA and history of hypersensitivity.

Warnings and Precautions

Monitor renal function, including blood urea nitrogen and serum creatinine, and urine output in patients at risk of developing acute renal failure. Hyperproteinemia, increased serum osmolarity, and hyponatremia may occur in patients receiving PANZYGA. Aseptic meningitis syndrome may occur in patients receiving PANZYGA, especially with high doses or rapid infusion. Hemolysis that is either intravascular or due to enhanced red blood cell sequestration can develop subsequent to PANZYGA treatments. Risk factors for hemolysis include high doses and non-O-blood group. Closely monitor patients for hemolysis and hemolytic anemia. Monitor patients for pulmonary adverse reactions (transfusion-related acute lung injury [TRALI]). Monitor blood pressure prior to, during, and following PANZYGA infusion. Carefully consider the relative risks and benefits before prescribing the high dose regimen (for cITP) in patients at increased risk of volume overload. PANZYGA is made from human plasma and may contain infectious agents, e.g. viruses and theoretically, the Creutzfeldt-Jakob disease agent.

Adverse Reactions

PI – The most common adverse reactions (≥5% study subjects) were headache, nausea, fever, fatigue, and abdominal pain.

cITP in adults – The most common adverse reactions (≥5% study subjects) were headache, fever, nausea, vomiting, dizziness, and anemia.

CIDP in adults – The most common adverse reactions reported in greater than 5% of subjects were: headache, fever, dermatitis, and blood pressure increase. The risk information provided here is not comprehensive; see full Prescribing Information and Boxed Warning for PANZYGA.

Please click [here](#) for Full Prescribing Information, including BOXED WARNING.